arXiv:1302.0353v1 [gr-qc] 2 Feb 2013

Hawking radiation and total entropy change as tunneling

Chikun Ding*

Department of Physics and Information Engineering, Hunan University of Humanities, Science and Technology, Loudi, Hunan 417000, P. R. China

Abstract

In the frame of Hamilton-Jacobi method, the back-reactions of the radiating particles together with the total entropy change of the whole system are investigated. The emission probability from this processing is found to be equivalent to the null geodesic method. However the total entropy of the system, particles and black hole, is increased yet very little, which is a little different. This method can be applied to any sort of horizons and particles without a specific choice of (regular-across-the-horizon) coordinates. Its physical meaning is more obvious: the negative energy one of a virtual particle pair is absorbed by the black hole, resulting in the temperature, electric potential and angular velocity increase; then the black hole amount of heat, electric charge and angular momentum can spontaneously transfer to the positive energy particle; when obtaining enough energy, it can escape away to infinity, visible to distant observers.

PACS numbers: 04.70.Dy, 04.62.+v

---

*Electronic address: dingchikun@163.com

I. INTRODUCTION

Using the quantum field theory in curved spacetime, Hawking [1] found that the collapsing black hole will lead, at late times, to a radiation of particles in all modes of the quantum field, with characteristic thermal spectrum at a temperature $1/8\pi M$. It is generally believed that the pair productions occur inside and outside the horizon of the black hole and tunnel across the horizon. In the late time, with knowledge of Feyman's [2] path integral method in the quantum mechanics, he found [3] that the probability of emission particles from the past horizon is not the same as the probability of absorption into the future horizon. The ratio for the Schwarzschild black hole between them is of the form $\Gamma_{out} = e^{-E/8\pi M}\Gamma_{in}$; for Reissner-Nordström black hole is $\Gamma_{out} = e^{-(E-qV_+)/T_R}\Gamma_{in}$; for Kerr black hole is $\Gamma_{out} = e^{-(E-m\Omega_+)/T_K}\Gamma_{in}$. These discovers have excited a lot of interest [4–6, 8–13, 15–32].

Being enlightened by path-integral method, K. Srinivasan *et al* [4–6] used Landau's [7] complex paths method to deduce radiance without using the Kruskal extension. They treated the radiance as tunnelling across the singularity and the wave functions as semiclassical approximation modes $\exp[\frac{i}{\hbar}I(r,t)]$, where I is the classical action function which can be expanded by $\hbar/i$. To the lowest order, I satisfies the relativistic Hamilton-Jacobi equation which gives a solution $I_\pm = -Et \pm W(r) + J(x^i)$, where "+" is of outgoing particles and "$-$" of incoming particles. I has a pole at horizon $r = r_+$ and the probabilities of the particles $\Gamma_{out} \sim e^{-2\mathrm{Im}I_+}$, $\Gamma_{in} \sim e^{-2\mathrm{Im}I_-}$.

This complex-path method has been known as Hamilton-Jacobi method after developed by Angheben *et al* [8] and man *et al* [9, 10]. To ensure the probability is normalized, they used the boundary conditions for incoming particles which fall behind the horizon along classically permitted trajectories, i.e. $I = -Et + W(r) + J(x^i) + K$, where K is a complex normalizing constant. So the total probability is

$$\Gamma = \Gamma_{out} \sim e^{-2[\mathrm{Im}I_+ - \mathrm{Im}I_-]}. \tag{1.1}$$

However this method can only be used under the condition that the background spacetime is considered fixed in which the energy conservation is not enforced during the emission process. Some efforts on extension this method to dynamic geometry have been done [11, 12]. But these generalizations have some crudeness.

Using usual thermodynamic way, we divide the emission process into many infinite small segments, every one of which can be treated as a quasi-static process and, the background spacetime can be treated as fixed, there exists equilibrium temperature. Thus in every segment we can use Hamilton-Jacobi method to handle. In different segment the instantaneous event horizon is different. We obtain each action I_i in every tiny time piece after the particle tunneled across the instantaneous horizon. To get the last action I, the change ΔI_i between the instantaneous actions should be considered. Then $I = \sum \Delta I_i \sim \int dI$. After integrating over the whole process, we obtain the thermal spectrum incorporating the effects of the back-reaction on the background spacetime, which is the same as that obtained by the null geodesic method proposed in [13].

There are two different approaches that are used to model tunneling process. The first method developed was the null geodesic method used by Parikh and Wilczek[13]. Another one is the Hamilton-Jacobi method. There have a couple of unpleasant features in the null geodesic method: (i) it strongly relies on a very specific choice of (regular-across-the-horizon) coordinates, and (ii) it turns upside down the relationship between Hawking's radiation and back-reaction [14]. The Hamilton-Jacobi method can cope with both issues. Therefore our method can be applied to any sort of horizon and particles.

We also study the change of total entropy of the system including black hole and radiating particles by investigating that where the particle energy comes from. Then we know why the entropy change of the black hole can be obtain by probing its radiating particles. Our result is that the difference of total entropy ΔS is very small but nonzero, which has some difference from ref. [13] in which the difference of total entropy $\Delta S = 0$. As in ref. [15], the authors argue that the null geodesic method is only suitable for the reversible process and the factual emission process is irreversible which is possible to lose information. Here we argue that Hamilton-Jacobi method can be suitable for the irreversible process and there are very few information lost in the emitting process.

Our paper is outlined as follows. Section II is for neutral scalar particles radiation from Reissner-Nordström black hole and the entropy change of the black hole and radiating particles.. Section III is devoted for charged scalar particles radiating from Reissner-Nordström black hole. Section IV is used discussing the charged scalar particles radiating from Kerr-Newman black hole. In section V, charged Dirac particles radiation from Kerr-Newman black hole is investigated. Section VI is a summary.

II. HAWKING RADIATION FOR REISSNER-NORDSTRÖM BLACK HOLE FROM NEUTRAL SCALAR PARTICLES RADIATION

The line element for the charged Reissner-Nordstrom back hole is described by

$$ds^2 = -\frac{\Delta(r)}{r^2}dt^2 + \frac{r^2}{\Delta(r)}dr^2 + r^2(d\theta^2 + \sin^2\theta d\varphi^2), \tag{2.1}$$

where $\Delta(r) = r^2 - 2Mr + Q^2$. When a neutral particle with the energy E tunnels out across the horizon, the black hole mass and electric charge M would be decreased to $M - E$ due to energy conservation, that is to say, the background spacetime is affected by the back-reaction of the emitting particle, $g^{\mu\nu}(r(M)) \to g^{\mu\nu}(r(M-E))$. However, because of quantum uncertainty principle, it seems too crude that the approximation of a discontinuous jump from M to $M-E$ for the black hole mass. Rather, it will require a "gradual" transition (relative to whatever time scale is characteristic of the radiation process). Therefore we divide this process into many infinite small segments, during ith one of which the particle obtains energy $\Delta\omega_i$, where $\Delta\omega_i = \omega_i - \omega_{i-1} \ll E$. These segments can be treat as many quasi-static processes and can be handled by Hamilton-Jacobi method.

A particle of instantaneous energy ω_i will effectively "sees" a spacetime metric of the form

$$ds^2 = -\frac{\Delta(r(M-\omega_i))}{r^2}dt^2 + \frac{r^2}{\Delta(r(M-\omega_i))}dr^2 + r^2(d\theta^2 + \sin^2\theta d\varphi^2). \tag{2.2}$$

In the following subsection, we use Hamilton-Jacobi method to study Hawking radiation incorporating back-reaction as tunneling, then the entropy change of the whole system of black hole and radiating particles in the next subsection.

A. The tunneling process

Now we divide the tunneling time t into infinite small pieces t_i and use Hamilton-Jacobi method to study these infinite small processes. The WKB approximation wave function is

$$\phi(t_i, r, \theta, \varphi) = \exp\left[\frac{i}{\hbar}I_i(t_i, r, \theta, \varphi) + I_1'(t_i, r, \theta, \varphi) + \mathcal{O}(\hbar)\right]. \tag{2.3}$$

The Klein-Gordon equation is

$$\frac{1}{\sqrt{-g(r(M-\omega_i))}}\partial_\mu\Big(\sqrt{-g(r(M-\omega_i))}\; g^{\mu\nu}(r(M-\omega_i))\partial_\nu\phi\Big) - \frac{u^2}{\hbar^2}\phi = 0. \tag{2.4}$$

Substituting Eq. (2.3) into (2.4), to leading order in $\hbar$, one get the following relativistic Hamilton-Jacobi equation

$$g^{\mu\nu}(r(M-\omega_i))(\partial_\mu I_i \partial_\nu I_i) + u^2 = 0, \tag{2.5}$$

where there exists a solution in the form

$$I_i = -\omega_i t_i + W_i(r) + J_i(\theta, \varphi) + K_i\ , \tag{2.6}$$

where K_i is a complex constant normalizing the action function. Substituting Eq. (2.2) into (2.5) yields

$$W_{i\pm}(r) \ = \ \pm\int \frac{r^2 dr}{\Delta(r(M-\omega_i))}\sqrt{\omega_i^2 - \frac{\Delta(r(M-\omega_i))}{r^2}(u^2 + g^{pk}J_pJ_k)}\ ,$$

where $J_p = \partial_p I_i$, $p = \theta, \varphi$. One solution of above corresponds to the scalar particles moving away from the black hole (i.e. "+" outgoing) and the other solution corresponds to particles moving toward the black hole (i.e. "−" incoming). Imaginary parts of the action can only come from the vicinity of the pole at the horizon. Integrating around the pole, the imaginary parts are

$$\mathrm{Im} I_{i\pm} = \mathrm{Im} W_{i\pm}(r) = \pm f(M-\omega_i)\,\omega_i + \mathrm{Im} K_i\ , \quad f(M-\omega_i) = \frac{\pi r_+'^2(M-\omega_i)}{2\sqrt{(M-\omega_i)^2 - Q^2}}, \tag{2.7}$$

where $r_+'(M-\omega_i) = M - \omega_i + \sqrt{(M-\omega_i)^2 - Q^2}$ is the instantaneous horizon. Therefore the action of the particle tunneled across the ith instantaneous horizon is

$$I_i = -\omega_i t_i + i f(M-\omega_i)\,\omega_i + K_i\ . \tag{2.8}$$

When the energy of the particle gradually approaches to E, its action is I. To obtain it, we should consider the change between the ith and $(i-1)$th instantaneous actions ΔI_i

$$\Delta I_i = -\Delta\omega_i\, t_i + i f(M-\omega_i)\,\Delta\omega_i + \Delta K_i\ . \tag{2.9}$$

Then

$$I = \sum \Delta I_i = \int_0^E -t d\omega + i f(M-\omega)\, d\omega + K\ . \tag{2.10}$$

At last the imaginary part of its action can be obtained by

$$\mathrm{Im} I = \int_0^E \frac{\pi r_+'^2}{2\sqrt{(M-\omega)^2 - Q^2}} d\omega + \mathrm{Im} K. \tag{2.11}$$

The imaginary parts of the action of tunneled particle are

$$\begin{aligned} \mathrm{Im} I_\pm &= \pm \int_0^E \frac{\pi(M-\omega+\sqrt{(M-\omega)^2-Q^2})^2}{2\sqrt{(M-\omega)^2-Q^2}} d\omega + \mathrm{Im} K \\ &= \pm\frac{\pi}{2}\big[2E(M-\frac{E}{2}) - (M-E)\sqrt{(M-E)^2-Q^2} + M\sqrt{M^2-Q^2}\ \big] + \mathrm{Im} K. \end{aligned} \tag{2.12}$$

Using Eq. (1.1), the emission probability is

$$\Gamma = e^{-2\pi\left(2E(M-\frac{E}{2}) - (M-E)\sqrt{(M-E)^2-Q^2} + M\sqrt{M^2-Q^2}\ \right)} = e^{\Delta S_{B-H}}, \tag{2.13}$$

where ΔS_{B-H} is the difference of Bekenstein-Hawking entropy of the black hole. This result is the same as in [13]. In the following subsection, we study the entropy change of the whole system and why the emission probability (2.13) is related to the entropy change of the black hole ΔS_{B-H}.

B. The total entropy change of the whole system

How does the particle obtain energy? If we consider the black hole and its radiation as a isolated system, then the particle energy can only come from absorbing black hole's amount of heat $\widetilde{Q}$. Because of vacuum fluctuation near the event horizon, at sometime the black hole temperature approaches to $T'(M-\omega_1)$, which is higher than the particle temperature $T'(M)$, so that amount of heat can spontaneously transfers from black hole to the particle.

In the first segment, particle energy $0 \to \omega_1$, the black hole entropy decreases,

$$\Delta S'_1 = -\widetilde{Q}_1/T'(M-\omega_1), \tag{2.14}$$

where S' is the black hole entropy, $T'(M-\omega_1) = \frac{1}{2\pi}\sqrt{(M-\omega_1)^2 - Q^2}/r'^2_+(M-\omega_1)$. The particle entropy S'' increases,

$$\Delta S''_1 = \widetilde{Q}_1/T'(M), \tag{2.15}$$

and the entropy of the system increase

$$\Delta S_1 = \Delta S''_1 + \Delta S'_1 = \widetilde{Q}_1/T'(M) - \widetilde{Q}_1/T'(M-\omega_1) > 0. \tag{2.16}$$

It shows that the radiating process is irreversible and ω_1 is

$$\Delta\omega_1 = \omega_1 - 0 = -T'(M-\omega_1)\Delta S'_1\ . \tag{2.17}$$

It is easy to see that we can really obtain the entropy change of the black hole by probing its radiating particle energy and, the emission probability (2.13) is really related to the entropy change of the black hole ΔS_{B-H}. At the end of the first segment, the particle temperature approached to $T'(M-\omega_1)$ and, for the black hole, it approached to $T'(M-\omega_2)$.

In the second segment, particle energy $\omega_1 \to \omega_2$, the particle absorbs heat $\Delta\omega_2 = \omega_2 - \omega_1 = -T'(M-\omega_2)\Delta S'_2$, and the increase of the system entropy is

$$\Delta S_2 = \Delta S''_2 + \Delta S'_2 = \Delta\omega_2/T'(M-\omega_1) - \Delta\omega_2/T'(M-\omega_2). \tag{2.18}$$

When the energy of the particle has approached to E, the black hole temperature is $T(M-E)$, and the total increase of the system entropy is

$$\Delta S = \Delta S_1 + \Delta S_2 + \cdots = \Delta\omega/T'(M) - \Delta\omega/T'(M-E) < \Delta\omega/T'(M) \tag{2.19}$$

under condition that $\Delta\omega_1 = \Delta\omega_2 = \cdots = \Delta\omega$. Due to $\Delta\omega \ll E$, the total increase of the system entropy is very small but nonzero. It is the same as in ref. [33], but has some difference from ref. [13] in which $\Delta S = 0$ (*In the limit the emitted particle carries away the entire mass and charge of the black hole, there are exp*(S_{B-H}) *states in total and the probability of findings a shell containing all of the mass of the black hole is proportional to exp*$(-S_{B-H})$).

III. HAWKING RADIATION FOR REISSNER-NORDSTRÖM BLACK HOLE FROM CHARGED SCALAR PARTICLES RADIATION

A particle of instantaneous energy ω_i and charge q'_i will effectively sees a spacetime metric of the form

$$ds^2 = -\frac{\Delta(r(M-\omega_i, Q-q'_i))}{r^2}dt^2 + \frac{r^2}{\Delta(r(M-\omega_i, Q-q'_i))}dr^2 + r^2(d\theta^2 + \sin^2\theta d\varphi^2). \tag{3.1}$$

The charged Klein-Gordon equation is

$$\frac{(\partial_\mu - iq'_iA_\mu)\Big(\sqrt{-g(r(M-\omega_i, Q-q'_i))}\; g^{\mu\nu}(r(M-\omega_i, Q-q'_i))(\partial_\nu - iq'_iA_\nu)\phi\Big)}{\sqrt{-g(r(M-\omega_i, Q-q'_i))}} - \frac{u^2}{\hbar^2}\phi = 0, \tag{3.2}$$

where $A_\mu = (-(Q-q'_i)/r, 0, 0, 0)$ is the potential of the electromagnetic field of the background spacetime. Substituting Eq. (2.3) into (3.2), to leading order in $\hbar$, one get the following relativistic Hamilton-Jacobi equation

$$g^{\mu\nu}(r(M-\omega_i, Q-q'_i))(\partial_\mu I_i \partial_\nu I_i + q'^2_i A_\mu A_\nu - 2q'_iA_\mu\partial_\nu I_i) + u^2 = 0. \tag{3.3}$$

Substituting Eq. (3.1) into (3.3) yields

$$W_{i\pm}(r) \;=\; \pm\int \frac{r^2dr}{\Delta(r(M-\omega_i, Q-q'_i))}\sqrt{\big[\omega_i - \frac{q'_i(Q-q'_i)}{r}\big]^2 - \frac{\Delta(r(M-\omega_i, Q-q'_i))}{r^2}(u^2 + g^{pk}J_pJ_k)}\;.$$

Integrating around the pole, the imaginary parts are

$$\mathrm{Im}I_{i\pm} = \mathrm{Im}W_{i\pm}(r) = \pm\frac{\pi r'^2_+(M-\omega_i,\ Q-q'_i)}{2\sqrt{(M-\omega_i)^2-(Q-q'_i)^2}}\Big(\omega_i - q'_iV'_+(M-\omega_i,\ Q-q'_i)\Big) + \mathrm{Im}K, \tag{3.4}$$

where $V'_+(M-\omega_i,\ Q-q'_i) = (Q-q'_i)/r'_+(M-\omega_i,\ Q-q'_i)$ is the instantaneous electric potential near the outer horizon, and $r'_+(M-\omega_i,\ Q-q'_i) = M-\omega_i + \sqrt{(M-\omega_i)^2-(Q-q'_i)^2}$ is instantaneous the horizon. The action of the particle tunneled across the ith instantaneous horizon is

$$I_i = -\omega_i t_i + i\frac{\pi r'^2_+}{2\sqrt{(M-\omega_i)^2-(Q-q'_i)^2}}(\omega_i - q'_iV'_+) + K_i\ , \tag{3.5}$$

and the change between the ith and $(i-1)$th instantaneous actions ΔI_i

$$\Delta I_i = -\Delta\omega_i\ t_i + i\frac{\pi r'^2_+}{2\sqrt{(M-\omega_i)^2-(Q-q'_i)^2}}(\Delta\omega_i - \Delta q'_iV'_+) + \Delta K_i\ , \tag{3.6}$$

where $\Delta q'_i = q'_i - q'_{i-1}$. When the energy of the particle gradually approaches to E, its action is I

$$I = \sum \Delta I_i = \int_0^E -td\omega + i\int_{(0,0)}^{(E,q)} \frac{\pi r'^2_+}{2\sqrt{(M-\omega_i)^2 - (Q-q'_i)^2}}(d\omega - dq'V'_+) + K \ . \tag{3.7}$$

Using Eq. (3.7), the imaginary parts of the action of tunneled particle are

$$\begin{aligned}\mathrm{Im}I_\pm &= \pm\int_{(0,0)}^{(E,q)} \frac{\pi(M-\omega+\sqrt{(M-\omega)^2-(Q-q')^2})^2}{2\sqrt{(M-\omega)^2-(Q-q')^2}}(d\omega - V'_+dq') + \mathrm{Im}K \\ &= \pm\frac{\pi}{2}\big[2E(M-\frac{E}{2}) - (M-E)\sqrt{(M-E)^2-(Q-q)^2} \\ &\quad + M\sqrt{M^2-(Q-q)^2} - Qq + \frac{1}{2}q^2\ \big] + \mathrm{Im}K. \end{aligned} \tag{3.8}$$

Using Eq. (1.1), the emission probability is

$$\Gamma = e^{-2\pi\left(2E(M-\frac{E}{2})-(M-E)\sqrt{(M-E)^2-(Q-q)^2}+M\sqrt{M^2-(Q-q)^2}-Qq+\frac{1}{2}q^2\ \right)} = e^{\Delta S_{B-H}}, \tag{3.9}$$

which is the same as in [20].

Now turn to the acquisition of particle energy. At sometime, the black hole temperature rises from $T'(M,\ Q)$ to $T'(M-\omega_1,\ Q-q'_1)$, its horizon shrinks from $r'_+(M,\ Q)$ to $r'_+(M-\omega_1,\ Q-q'_1)$, which lead to increase of the electric potential of the black hole, $V'_+(M,\ Q) \to V'_+(M-\omega_1,\ Q-q'_1)$, where $T'(M-\omega_1,\ Q-q'_1) = \frac{1}{2\pi}\sqrt{(M-\omega_1)^2-(Q-q'_1)^2}/r'^2_+(M-\omega_1,\ Q-q'_1)$. Then the amount of heat and electric charge flow to the particle, which will lead to another further increase of black hole temperature and electric potential. The particle energy comes from two ways, absorbing black hole's internal energy and electric potential energy $q'_iV'_+$. Energy ω_i is

$$\Delta\omega_i = \omega_i - \omega_{i-1} = -T'(M-\omega_i)\Delta S'_i + V'_+\Delta q'_i \ , \tag{3.10}$$

where $\Delta q'_i = q'_i - q'_{i-1}$.

IV. HAWKING RADIATION FOR KERR-NEWMAN BLACK HOLE FROM CHARGED SCALAR PARTICLES RADIATION

The "no hair" theorem stated that all information about the collapsing body was lost from the outside region apart from three conserved quantities: the mass, the angular momentum and the electric charge, the final state of most of collapsing star is Kerr-Newman black hole.

In the Boyer-Lindquist coordinate, its line element in four dimensional spacetime is described by

$$\begin{aligned} ds^2 = &-\left(1-\frac{2Mr}{\rho^2}\right)dt^2 - \frac{4Mra\sin^2\theta}{\rho^2}dtd\varphi + \frac{\rho^2}{\triangle}dr^2 \\ &+\rho^2 d\theta^2 + \left(r^2+a^2+\frac{2Mra^2\sin^2\theta}{\rho^2}\right)\sin^2\theta d\varphi^2, \end{aligned} \tag{4.1}$$

with

$$\begin{aligned} &\rho^2 = r^2 + a^2\cos^2\theta, \quad \triangle = r^2 - 2Mr + a^2 + Q^2 = (r-r_+)(r-r_-) \\ &r_+ = M + \sqrt{M^2-a^2-Q^2}, \quad r_- = M - \sqrt{M^2-a^2-Q^2}, \end{aligned} \tag{4.2}$$

where M is the mass of the black hole and $a = J/M$ is the angular momentum parameter; r_- and r_+ are the inner and event horizons; its electromagnetic field potential is $A_\mu = (-Qr/\rho^2,\ 0,\ 0,\ Qra\sin^2\theta/\rho^2)$. When a particle with the energy E, electric charge q and angular momentum j tunnels out across the horizon, the black hole mass, angular momentum, and electric charge $M,\ Q,\ J$ would be decreased to $M-E,\ Q-q,\ J-j$ due to energy, electric charge and angular momentum conservation. We also divide this process into many infinite small segments, during the ith one of which the particle obtains energy $\Delta\omega_i$, angular momentum $\Delta j_i'$, where $\Delta j_i' = j_i' - j_{i-1}'$. These segments can be treat as many quasi-static processes and can be handled by Hamilton-Jacobi method.

A particle of instantaneous energy ω_i, charge q_i' and angular momentum j_i' will effectively sees a spacetime metric of the form

$$\begin{aligned} ds^2 = &-\left(1-\frac{2(M-\omega_i)r}{\tilde{\rho}^2}\right)dt^2 - \frac{4(M-\omega_i)r\tilde{a}_i\sin^2\theta}{\tilde{\rho}^2}dtd\varphi + \frac{\tilde{\rho}^2}{\triangle(r(M-\omega_i, Q-q_i', J-j_i'))}dr^2 \\ &+\tilde{\rho}^2 d\theta^2 + \left(r^2+\tilde{a}_i^2+\frac{2(M-\omega_i)r\tilde{a}_i^2\sin^2\theta}{\tilde{\rho}^2}\right)\sin^2\theta d\varphi^2, \end{aligned} \tag{4.3}$$

where $\tilde{a}_i = (J-j_i')/(M-\omega_i)$, $\tilde{\rho}^2 = r^2 + \tilde{a}_i^2\cos^2\theta$.

We divide tunneling time t, rotating angle φ into infinite small pieces $t_i,\ \varphi_i$ and use Hamilton-Jacobi method. Due to the energy and angular momentum conservation, its instantaneously action function must be of the form

$$I_i = -\omega_i t_i + W_i(r) + j_i'\varphi_i + \tilde{J}_i(\theta) + K_i. \tag{4.4}$$

Taking the line element (4.3) into Hamilton-Jacobi equation (3.3), one find

$$\begin{aligned} &\triangle^2 W_i'^2(r) + \left[\triangle u^2 r^2 - \tilde{a}_i^2\widetilde{j_i'}^2 + 4(M-\omega_i)r\tilde{a}_i\widetilde{j_i'}\widetilde{\omega}_i - (r^2+\tilde{a}_i^2)^2\widetilde{\omega}_i^2\right] + \triangle\lambda = 0, \\ &\lambda = \left[\tilde{a}_i^2\sin^2\theta\widetilde{\omega}_i^2 + \tilde{J}_i'^2(\theta) + \frac{\widetilde{j_i'}^2}{\sin^2\theta} + u^2\tilde{a}_i^2\cos^2\theta\right], \end{aligned} \tag{4.5}$$

where

$$\widetilde{\omega}_i = \omega_i - \frac{q_i'(Q-q_i')r}{\tilde{\rho}^2}, \quad \widetilde{j_i'} = j' - \frac{q_i'(Q-q_i')r}{\tilde{\rho}^2}\tilde{a}_i \sin^2\theta. \tag{4.6}$$

Then the imaginary parts of the action function are

$$\begin{aligned}
\mathrm{Im}I_{i\pm} &= \pm\mathrm{Im}\left[\int dr \frac{1}{\triangle}\sqrt{(r^2+\tilde{a}_i^2)^2\big(\widetilde{\omega}_i - \frac{\widetilde{j_i'}\tilde{a}_i}{r^2+\tilde{a}_i^2}\big)^2 - \triangle(u^2r^2+\lambda-2\widetilde{j_i'}\tilde{a}_i\widetilde{\omega}_i)}\right] + \mathrm{Im}K_i \\
&= \pi\frac{r_+'^2+\tilde{a}_i^2}{2(r_+'-M+\omega_i)}\left(\widetilde{\omega}_{i+} - \frac{\widetilde{j_{i+}'}\tilde{a}_i}{r_+'^2+\tilde{a}_i^2}\right) + \mathrm{Im}K_i \\
&= \pi\frac{r_+'^2+\tilde{a}_i^2}{2(r_+'-M+\omega_i)}\left(\omega_i - \frac{q_i'(Q-q_i')r_+'}{r_+'^2+\tilde{a}_i^2} - \frac{j_i'\tilde{a}_i}{r_+'^2+\tilde{a}_i^2}\right) + \mathrm{Im}K_i \\
&= \pi\frac{r_+'^2+\tilde{a}_i^2}{2(r_+'-M+\omega_i)}\left(\omega_i - q_i'V_+' - j_i'\Omega_+'\right) + \mathrm{Im}K_i,
\end{aligned} \tag{4.7}$$

where $r_+'(M-\omega_i,\ Q-q_i',\ J-j_i') = M-\omega_i+\sqrt{(M-\omega_i)^2-(Q-q_i')^2-\tilde{a}_i^2}$, $V_+'(M-\omega_i,\ Q-q_i',\ J-j_i') = (Q-q_i')r_+'/(r_+'^2+\tilde{a}_i^2)$ is the electromagnetic potential on the horizon, $\Omega_+'(M-\omega_i,\ Q-q_i',\ J-j_i') = \tilde{a}_i/(r_+'^2+\tilde{a}_i^2)$ is the dragging velocity of the horizon.

When the energy, charge and angular momentum of the particle gradually approaches to $E,\ q,\ j$, the imaginary part of its action can be obtained by

$$\mathrm{Im}I = \int_{(0,0,0)}^{(E,q,j)} \pi\frac{r_+'^2+\tilde{a}_i^2}{2(r_+'-M+\omega)}(d\omega - V_+'dq' - \Omega_+'dj') + \mathrm{Im}K. \tag{4.8}$$

Using Eq. (4.8), the imaginary parts of the action of tunnelled particle are

$$\begin{aligned}
\mathrm{Im}I_\pm &= \pm\frac{\pi}{2}\big[2E(M-\frac{E}{2}) - (M-E)\sqrt{(M-E)^2-(Q-q)^2-\tilde{a}^2} \\
&\quad + M\sqrt{M^2-(Q-q)^2-a^2} - Qq + \frac{1}{2}q^2 + \tilde{a}^2 - a^2\ \big] + \mathrm{Im}K,
\end{aligned} \tag{4.9}$$

where $\tilde{a} = (J-j)/(M-E)$.

Using Eq. (1.1), the emission probability is

$$\Gamma = e^{-2\pi\left(2E(M-\frac{E}{2})-(M-E)\sqrt{(M-E)^2-(Q-q)^2-\tilde{a}^2}+M\sqrt{M^2-(Q-q)^2-a^2}-Qq+\frac{1}{2}q^2+\tilde{a}^2-a^2\ \right)} = e^{\Delta S_{B-H}}.$$

If set $j = Ea$ for special case, then $\tilde{a} = a$, so this result is the same as in [21, 22].

Now turn to the acquisition of particle energy near the horizon of Kerr-Newman black hole. At sometime, the black hole temperature rises from $T'(M,\ Q,\ J)$ to $T'(M-\omega_1,\ Q-q_1',\ J-j_1')$, its horizon shrinks from $r_+'(M,\ Q,\ J)$ to $r_+'(M-\omega_1,\ Q-q_1',\ J-j_1')$, which lead to increase of the electric potential and the angular velocity of the black hole, $V_+'(M,\ Q,\ J) \to V_+'(M-\omega_1,\ Q-q_1',\ J-j_1')$, $\Omega_+'(M,\ Q,\ J) \to \Omega_+'(M-\omega_1,\ Q-q_1',\ J-j_1')$, where $T'(M-\omega_1,\ Q-q_1',\ J-j_1') =$

$\frac{1}{2\pi}\sqrt{(M-\omega_1)^2-(Q-q_1')^2-\tilde{a}_1^2}/(r_+'^2+\tilde{a}_1^2)$. Then the amount of heat, electric charge and angular momentum flow to the particle, which will lead to another further increase of black hole temperature, electric potential and angular velocity. The particle energy comes from three ways, absorbing black hole's internal energy, electric potential energy $q_i'V_+'$ and rotational kinetic energy $\Omega_+'\Delta j_i'$

$$\Delta\omega_i = -T'(M-\omega_i)\Delta S_i' + V_+'\Delta q_i' + \Omega_+'\Delta j_i' \ . \tag{4.10}$$

V. HAWKING RADIATION FOR KERR-NEWMAN BLACK HOLE FROM CHARGED DIRAC PARTICLES TUNNELLING

In this section, we extend the Hamilton-Jacobi method to Dirac field. The charged Dirac equation is

$$\left[\gamma^\alpha e_\alpha^\mu(\partial_\mu + \Gamma_\mu - iq_i'A_\mu) + \frac{u}{\hbar}\right]\psi = 0, \tag{5.1}$$

with

$$\Gamma_\mu = \frac{1}{8}[\gamma^a, \gamma^b]e_a^\nu e_{b\nu;\mu},$$

where γ^a are the Dirac matrices and e_a^μ is the inverse tetrad defined by $\{e_a^\mu\gamma^a, \quad e_b^\nu\gamma^b\} = 2g^{\mu\nu}\times 1$. For the Kerr-Newman metrics in Boyer-Lindquist coordinate (4.3), the nonzero tetrad elements can be taken as

$$e_0^{t_i} = \frac{1}{\sqrt{1-\frac{2(M-\omega_i)r}{\tilde{\rho}^2}}}, \quad e_3^{t_i} = -\frac{2(M-\omega_i)r\tilde{a}_i\sin\theta}{\tilde{\rho}\sqrt{\triangle^2-\tilde{a}_i^2\sin^2\theta\triangle}},$$

$$e_1^r = \frac{\sqrt{\triangle}}{\tilde{\rho}}, \qquad e_2^\theta = \frac{1}{\tilde{\rho}}, \quad e_3^{\varphi_i} = \frac{\sqrt{\triangle-\tilde{a}_i^2\sin^2\theta}}{\tilde{\rho}\sin\theta\sqrt{\triangle}},$$

where $\triangle = r^2+\tilde{a}_i^2-2(M-\omega_i)r+(Q-q_i')^2$. We employ the following ansatz for the Dirac field

$$\psi_{i\uparrow} = \begin{pmatrix} A(t_i,r,\theta,\varphi_i)\xi_\uparrow \\ B(t_i,r,\theta,\varphi_i)\xi_\uparrow \end{pmatrix} \exp\big(\frac{i}{\hbar}I_{i\uparrow}(t_i,r,\theta,\varphi_i)\big) = \begin{pmatrix} A(t_i,r,\theta,\varphi_i) \\ 0 \\ B(t_i,r,\theta,\varphi_i) \\ 0 \end{pmatrix} \exp\big(\frac{i}{\hbar}I_{i\uparrow}(t_i,r,\theta,\varphi_i)\big),$$

$$\psi_{i\downarrow} = \begin{pmatrix} C(t_i,r,\theta,\varphi_i)\xi_\downarrow \\ D(t_i,r,\theta,\varphi_i)\xi_\downarrow \end{pmatrix} \exp\big(\frac{i}{\hbar}I_{i\downarrow}(t_i,r,\theta,\varphi_i)\big) = \begin{pmatrix} 0 \\ C(t_i,r,\theta,\varphi_i) \\ 0 \\ D(t_i,r,\theta,\varphi_i) \end{pmatrix} \exp\big(\frac{i}{\hbar}I_{i\downarrow}(t_i,r,\theta,\varphi_i)\big),$$

$$\tag{5.2}$$

where "↑" and "↓" represent the spin up and spin down cases, and $\xi_{\uparrow}$ and $\xi_{\downarrow}$ are the eigenvectors of σ^3. Inserting Eqs. (5.2), (5.2) into the Dirac equation (5.1) and employing $I_{i\uparrow} = -\omega_i t_i + W_i(r) + j'_i\varphi_i + \tilde{J}_i(\theta) + \mathrm{Im}K_i$, to the lowest order in $\hbar$ we obtain

$$-e_0^{t_i} A\Big(\omega_i - \frac{q'_i(Q-q'_i)r}{\tilde{\rho}^2}\Big) + e_1^r B W'_i(r) + uA = 0, \tag{5.3}$$

$$e_0^{t_i} B\Big(\omega_i - \frac{q'_i(Q-q'_i)r}{\tilde{\rho}^2}\Big) - e_1^r A W'_i(r) + uB = 0, \tag{5.4}$$

$$B\left[-ie_3^{t_i}\Big(\omega_i - \frac{q'_i(Q-q'_i)r}{\tilde{\rho}^2}\Big) + e_2^{\theta}\tilde{J}_i{}'(\theta) + ie_3^{\varphi_i}\Big(j'_i - \frac{q'_i(Q-q'_i)r}{\tilde{\rho}^2}\tilde{a}_i\sin^2\theta\Big)\right] = 0, \tag{5.5}$$

$$-A\left[-ie_3^{t_i}\Big(\omega_i - \frac{q'_i(Q-q'_i)r}{\tilde{\rho}^2}\Big) + e_2^{\theta}\tilde{J}_i{}'(\theta) + ie_3^{\varphi_i}\Big(j'_i - \frac{q'_i(Q-q'_i)r}{\tilde{\rho}^2}\tilde{a}_i\sin^2\theta\Big)\right] = 0, \tag{5.6}$$

where we consider only the positive frequency contributions without loss of generality. From above four equations, we can obtain

$$\begin{aligned}
&\frac{\triangle}{\tilde{\rho}^2}W_i'^2(r) + u^2 - \frac{\Big(\omega_i - \frac{q'_i(Q-q'_i)r}{\tilde{\rho}^2}\Big)^2}{\rho^2\triangle}\left[\frac{4(M-\omega_i)^2 r^2\tilde{a}_i^2\sin^2\theta}{\triangle - \tilde{a}_i^2\sin^2\theta} + (r^2+\tilde{a}_i^2)^2 - \triangle\tilde{a}_i^2\sin^2\theta\right] = 0,\\
&\frac{1}{\tilde{\rho}^2}\left[\tilde{J}_i{}'^2(\theta) + \frac{\Big(j'_i - \frac{q'_i(Q-q'_i)r}{\tilde{\rho}^2}\tilde{a}_i\sin^2\theta\Big)^2(\triangle - \tilde{a}_i^2\sin^2\theta)}{\triangle\sin^2\theta} + \frac{4(M-\omega_i)^2 r^2\tilde{a}_i^2\Big(\omega_i - \frac{q'_i(Q-q'_i)r}{\tilde{\rho}^2}\Big)^2\sin^2\theta}{\triangle(\triangle - \tilde{a}_i^2\sin^2\theta)}\right.\\
&\left.+2\Big(j'_i - \frac{q'_i(Q-q'_i)r}{\tilde{\rho}^2}\tilde{a}_i\sin^2\theta\Big)\Big(\omega_i - \frac{q'_i(Q-q'_i)r}{\tilde{\rho}^2}\Big)\frac{2(M-\omega_i)r\tilde{a}_i}{\triangle}\right] = 0.
\end{aligned} \tag{5.7}$$

After calculating the above two equations, we find

$$\begin{aligned}
&\triangle^2 W_i'^2(r) - (r^2+\tilde{a}_i^2)^2\widetilde{\omega}_i^2 - \tilde{a}_i^2\widetilde{j'_i}^2 + 4(M-\omega_i)r\tilde{a}_i\widetilde{j'_i}\widetilde{\omega}_i\\
&+\triangle\left[\tilde{J}_i{}'^2(\theta) + \frac{\widetilde{j'_i}^2}{\sin^2\theta} + u^2r^2 + u^2\tilde{a}_i^2\cos^2\theta - \tilde{a}_i^2\sin^2\theta\,\widetilde{\omega}_i^2\right] = 0,
\end{aligned}$$

which is the same as the Eq. (4.5), so the Hawking radiation is again recovered. The result can be interpreted that the black holes radiate different spin weight of particles at the same temperature.

VI. SUMMARY

In the Hamilton-Jacobi framework, we have naturally discussed Reissner-Nordström and Kerr-Newman black holes' radiance with back-reaction from neutral scalar, charged scalar and Dirac particles radiation. Hamilton-Jacobi method can only be used to modeling the case that the background geometry is considered fixed. To handle the back-reaction of the radiating particles, we should first divide the radiation time into a series of infinite small pieces in which the small process can be a quasi-static one. Then use Hamilton-Jacobi method to obtain the instantaneous action of the particle tunneled across the instantaneous black hole horizon. To get the last action, we should find the changes between the ith and $(i-1)$th instantaneous action and sum them. The last result is the same as that obtained via the null geodesic method.

The physical meaning is obvious in this processing. Due to vacuum fluctuation near horizon, a virtual particle pair is created. The negative energy particle is absorbed by the black hole, resulting in the black hole mass decrease, while temperature, electric potential and angular velocity increase. Then the black hole amount of heat, electric charge and angular momentum can spontaneously transfer to the positive energy particle. This process also results in the black hole temperature, electric potential and angular velocity further increase and further transfer. When the particle obtains enough energy, it can escape away to infinity, visible to distant observers.

We have also studied the change of total entropy of the system including black hole and radiating particles and, answered that why the entropy change of the black hole can be obtain by probing its radiating particles. Our result is that the difference of total entropy ΔS is very small.

Acknowledgments

This work was supported by the National Natural Science Foundation of China under No. 11247013; Hunan Provincial NSFC No. 11JJ3014, the Scientific Research Fund of Hunan Provincial Education Department No. 11B067 and, the Foundation for the Author of Hunan Provincial Excellent Doctoral Dissertation No. YB2012B034; Aid program for Science and Technology Innovative Research Team in Higher Educational Institutions of Hunan Province.

---

[1] S. W. Hawking, Nature **248**, 30 (1974); Commun. Math. Phys. **43**, 199 (1975).

[2] R. P. Feyman, Rev. Mod. Phys. **20**, 327 (1948).

[3] J. B. Hartle and S. W. Hawking, Phys. Rev. D **13**, 2188 (1976).

[4] K. Srinivasan and T. Padmanabhan, Phys. Rev. D **60** 024007 (1999).

[5] S. Shankaranarayanan and K. Srinivasan and T. Padmanabhan, Mod. Phys. Lett. A **16** 571 (2001).

[6] S. Shankaranarayanan and T. Padmanabhan and K. Srinivasan, Class. Quantum Grav. **19** 2671 (2002).

[7] L. D. Landau and E. M. Lifshitz, *Quantum Mechanics (Nonrelativistic Theory)*, Course of Theoretical Physics, Volume 2 (Pergamon, New York, 1975).

[8] M. Angheben, M. Nadalini, L. Vanzo, and S. Zerbini, J. High Energy Phys. 0505 (2005) 014.

[9] R. Kerner and R. B. Mann, Phys. Rev. D**73** 104010 (2006).

[10] R. Kerner and R. B. Mann, Class. Quantum Grav. **25** 095014 (2008).

[11] A. J. M. Medved and E. C. Vagenas, Mod. Phys. Lett. A **20** 2449 (2005).

[12] S. Yang and D. Chen, Int. J. Theor. Phys. **46** 1747 (2007); D. Chen, X. Zu and S. Yang, Acta Phys. Polo. B **39** 1329 (2008).

[13] M. K. Parikh and F. Wilczek, Phys. Rev. Lett. **85** 5042 (2000).

[14] L. Vanzo, G. Acquaviva and R. Di Criscienzo, Class. Quantum Grav. **28** 183001 (2011).

[15] J. Zhang, Y. Hu and Z. Zhao, Mod. Phys. Lett. A **21**, 1865 (2006).

[16] P. Kraus, and Frank Wilczek, Mod. Phys. Lett. A **9**, 3713 (1994).

[17] P. Kraus and F. Wilczek, Nucl. Phys. B**433** 403 (1995).

[18] P. Kraus and F. Wilczek, Nucl. Phys. B**437** 231 (1995).

[19] B. K. Berger, D. M. Chitre, V. E. Moncrief and Y. Nutku, Phys. Rev. D**5** 2467 (1972).

[20] Jingyi Zhang, and Zheng Zhao, JHEP 10 (2005) 055.

[21] Jingyi Zhang, and Zheng Zhao, Phys. Lett. B**638** 110 (2006);

[22] Qing-Quan Jiang, Shuang-Qing Wu, and Xu Cai, Phys.Rev. D**73** 064003 (2006).

[23] M. K. Parikh, Phys. Lett. B **546** 189 (2002).

[24] M. K. Parikh, Int. J. Mod. Phys. D**13** 2351 (2004).

[25] M. K. Parikh, arXiv: hep-th/0402166.

[26] M. Arzano, A. Medved and E. Vagenas, J. High Energy Phys. 0509 (2005) 037.

[27] Shuang-Qing Wu, and Qing-Quan Jiang, J. High Energy Phys. 0603 (2006) 079.

[28] S. Q. Wu and X. Cai, Gen. Rel. Grav. **34** 1207 (2002).

[29] Shuang-Qing Wu and Mu-Lin Yan, Chin. Phys. Lett. **20** 1913 (2003).

[30] A. J. M. Medved and E. Vagenas, Mod. Phys. Lett. A**20** 2449 (2005).

[31] Bhramar Chatterjee, Amit Ghosh, P. Mitra, arXiv: 0704. 1746.

[32] P. Mitra, Phys. Lett. B**648** 240 (2007).

[33] W. H. Zurek, Phys. Rev. Lett. **49**, 1683-1686 (1982); D. N. Page, *ibid*, **50**, 1013 (1983).